BOUNDS ON THE DISTRIBUTION OF A SUM OF TWO RANDOM VARIABLES: Revisiting a problem of Kolmogorov with application to Individual Treatment Effects

Zhehao Zhang and Thomas S. Richardson University of Washington, Seattle Department of Statistics

May 15, 2024

Abstract

We revisit the following problem, proposed by Kolmogorov: given prescribed marginal distributions F and G for random variables X, Y respectively, characterize the set of compatible distribution functions for the sum Z = X + Y. Bounds on the distribution function for Z were given by Makarov (1982), and Frank et al. (1987), the latter using copula theory. However, though they obtain the same bounds, they make different assertions concerning their sharpness. In addition, their solutions leave some open problems in the case when the given marginal distribution functions are discontinuous. These issues have led to some confusion and erroneous statements in subsequent literature, which we correct.

Kolmogorov's problem is closely related to inferring possible distributions for individual treatment effects $Y_1 - Y_0$ given the marginal distributions of Y_1 and Y_0 ; the latter being identified from a randomized experiment. We use our new insights to sharpen and correct results due to Fan and Park (2010) concerning individual treatment effects, and to fill some other logical gaps.

1 Introduction

The question of the best possible bounds for the distribution function of the sum of two random variables whose individual distribution functions are fixed was originally raised by A.N. Kolmogorov and was first solved by Makarov (1982). Using copula theory (Sklar 1959), Frank et al. (1987) reframed this question and provided an elegant proof that the bounds were achievable in certain settings. Williamson and Downs (1990) generalized the bounds in Frank et al. (1987) to other arithmetic operations of two random variables including subtraction, multiplication and division and claimed these bounds are sharp. More recently, Fan and Park (2010) introduced the bounds into the context of causal inference and concluded sharp bounds on the distribution function of the additive treatment effect contrast (which corresponds to a difference between two random variables). The bounds proposed in Fan and Park (2010) have gained widespread traction in the literature on causal inference and econometrics, in works such as Chiba 2017, Huang et al. 2017, Lu et al. 2018 and Mullahy 2018.

In this paper, we revisit the connection between the result of Frank et al. (1987) and Kolmogorov's original question. We distinguish between bounds that are *achievable* and those that cannot be improved, and characterize the circumstances under which the bounds are achievable. We identify and address logical gaps in Williamson and Downs (1990) that have propagated to some of the later literature, leading to incorrect or imprecise statements. In particular, for the distribution function of the difference of two random variables, the lower bound proposed by Williamson and Downs is not sharp for measures that are not absolutely continuous with respect to Lebesgue measure. We also identify an unnecessary exclusion in the argument given by Williamson and Downs (1990). Building on Makarov (1982) and Frank et al. (1987), we formulate sharp bounds on the distribution function of the difference of two random variables with fixed marginals; we show how these differ from those appearing in previous literature. Finally, we apply the new bounds in the context of treatment effects and provide examples of how to calculate the bounds.

The paper is organized as follows. In Section 2, we review the proof of the best possible bounds proposed in Frank et al. (1987). In Section 3, we revisit the connection of Frank et al. (1987)'s result to Komogorov's question. We provide characterizations of when the bounds proposed in Frank et al. (1987) for the sum of two random variables are achievable. In Section 4, we state our new bounds for the difference of two random variables whose individual distribution functions are fixed. We discuss the implications of some of the results in Williamson and Downs (1990). In Section 5, we set up the problem of causal inference and revisit the bounds in Fan and Park (2010). We provide an example with discrete random variables in which the bounds are not sharp.

2 Bounds on the sum of two random variables

Given marginal distribution functions F, G for random variables X, Y respectively, Kolmogorov's question (restated here in terms of right-continuous distribution functions) is to find functions \underline{J} and \overline{J} such that for all $z \in \mathbb{R}$,

$$\underline{J}(z) = \inf P(X + Y \le z) \tag{1}$$

$$\overline{J}(z) = \sup P(X + Y \le z) \tag{2}$$

where the infimum and supremum are taken over all possible joint distribution functions H(x, y)having the marginal cdfs F(x) and G(y).

First, we review some existing results on probability distributions and copulas.

Definition 1. Let X be a random variable. The distribution function (or cumulative distribution function, cdf) F of X is defined to be $F(x) = P(X \le x)$ for $x \in \mathbb{R}$.

Note that under Definition 1, for any random variable X, $F(\cdot)$ is a right-continuous function. Frank et al. (1987) and Williamson and Downs (1990) used a left-continuous version of the definition of distribution functions where they replace $P(X \le x)$ with P(X < x).¹

Definition 2 (Embrechts and Hofert 2013). Let X be a random variable with distribution function F. The generalized inverse (also known as the quantile function) $F^{-1}: [0,1] \to \overline{\mathbb{R}} = [-\infty, \infty]$ of F is defined as:

$$F^{-1}(u) = \inf\{x \in \mathbb{R}, F(x) \ge u\}, u \in [0, 1]$$

with $\inf \emptyset = \infty$.

¹Some of the results we cited in this paper was originally defined in terms of the left-continuous functions $\tilde{F}(x) = P(X < x)$ and $\tilde{G}(x) = P(X < x)$ (which somewhat confusingly they also call "distribution functions"). To be consistent in notation, F, G in this paper always refer to the right-continuous distribution functions given by $F(x) = P(X \le x)$ and likewise for G and Y. We reserve \tilde{F}, \tilde{G} when we need to talk about the functions given by P(X < x) and P(Y < y), equivalently the left hand limits of F(x) and G(y).

Definition 3. A two dimensional copula is a mapping C from $[0,1]^2$ to [0,1] satisfying the conditions:

1.
$$C(a,0) = C(0,a) = 0$$
 and $C(a,1) = C(1,a) = a$, for all a in $[0,1]$;

2. $C(a_2, b_2) - C(a_1, b_2) - C(a_2, b_1) + C(a_1, b_1) \ge 0$ for all a_1, a_2, b_1, b_2 in [0, 1] such that $a_1 \le a_2, b_1 \le b_2$.

Proposition 4. According to the definition, any copula C is nondecreasing in each argument, and,

$$W(a,b) \le C(a,b) \le M(a,b) \tag{3}$$

where

$$W(a,b) = \max(a+b-1,0), \quad M(a,b) = \min(a,b).$$

The bounds W and M are known as Fréchet-Hoeffding copula bounds.

Theorem 5 (Sklar 1959). Consider a 2-dimensional cdf H with marginals F, G. There exists a copula C, such that

$$H(x, y) = C(F(x), G(y))$$

for all x, y in $[-\infty, \infty]$. If F, G are both continuous, then C is unique²; otherwise C is uniquely determined only on Ran $F \times \text{Ran } G$, where Ran F, Ran G denote respectively the range of the cdfs F and G.

See Sklar (1959), Embrechts and Hofert (2013), Schmidt (2007) for more discussion of general *n*-dimensional copulas and Sklar's Theorem.

Now we want to bound the cdf of the sum of two random variables using copulas. We reprise the argument given by Frank et al. (1987) which gives lower bounds on P(X + Y < z) and upper bounds on $P(X + Y \le z)$, and further establishes by construction that these are *achievable*.

Let *H* be a two-dimensional cumulative distribution function for random variables *X*, *Y* with marginals *F*, *G* respectively so that $H(x, y) = P(X \le x, Y \le y)$. By Sklar's theorem (Sklar 1959),

²Note that the condition here relates to the cdf F, G viewed as functions. We are not assuming that the random variables X, Y are (absolutely) continuous (with respect to Lebesgue measure). The latter is a sufficient but not necessary condition for F, G to be continuous.

there exists a copula C such that H(x, y) = C(F(x), G(y)). Note that the cdf for the sum of two random variables X, Y is fully characterized by their joint cdf H. Let Z := X + Y and J be the cdf for Z. Then for any $z \in [-\infty, \infty]$,

$$J(z) = \iint_{x+y \le z} dH(x,y). \tag{4}$$

For the copula C and marginal distribution functions F, G, let $\sigma_C(F, G)$ be the function defined by $\sigma_C(F, G)(-\infty) = 0, \sigma_C(F, G)(\infty) = 1$ and

$$\sigma_C(F,G)(z) = \iint_{x+y \le z} dC(F(x), G(y)), \quad \text{ for } -\infty < z < \infty$$

Since $H(x,y) = C(F(x),G(y)), \sigma_C(F,G)(z) = J(z)$ for all $z \in [-\infty,\infty]$. We let

$$\tau_C(F,G)(z) = \sup_{x+y=z} C(F(x), G(y)),$$
(5)

$$\rho_C(F,G)(z) = \inf_{x+y=z} \check{C}(F(x), G(y)), \tag{6}$$

where

$$\check{C}(a,b) = a + b - C(a,b).$$

Theorem 6 (Frank et al. Theorem 2.14). We have the following bounds for any copula C and arbitrary given distribution functions F, G and $z \in [-\infty, \infty]$:

$$\tau_W(F,G)(z) \le \tau_C(F,G)(z) \le \sigma_C(F,G)(z) \le \rho_C(F,G)(z) \le \rho_W(F,G)(z).^3$$

Note that τ_W, ρ_W are known functions that depend solely on the marginal cdfs F and G. We obtain the following bounds:

$$\tau_W(F,G)(z) \le \sigma_C(F,G)(z) \le \rho_W(F,G)(z) \tag{7}$$

³Proof of these bounds and visualizations can be found in Frank et al. (1987), Nelsen (2006). Here only σ_C can be interpreted as the probability that $Z \leq z$. Therefore, there are no explicit constructions establishing whether or not the bounds on τ_W , ρ_W can be achieved.

Here we will write the bounds τ_W, ρ_W explicitly.

$$\tau_W(F,G)(z) = \sup_{x+y=z} \max(F(x) + G(y) - 1, 0)$$
(8)

$$\rho_W(F,G)(z) = \inf_{x+y=z} \left(F(x) + G(y) - \max(F(x) + G(y) - 1, 0) \right) \tag{9}$$

$$= \inf_{x+y=z} \left(F(x) + G(y) + \min(-F(x) - G(y) + 1, 0) \right)$$
(10)

$$= \inf_{x+y=z} \min(1, F(x) + G(y))$$
(11)

$$= 1 + \inf_{x+y=z} \min(0, F(x) + G(y) - 1)$$
(12)

Theorem 6 establishes the validity of the bounds in Equation (8) and (12). Thereafter, we call the bounds $\tau_W(F,G)(z)$ and $\rho_W(F,G)(z)$ the Makarov bounds⁴.

3 Sharpness of the bounds

To investigate the tightness of the bounds, we first distinguish three notions of sharpness.

For two random variables X, Y with fixed marginals F, G, respectively, let $J(\cdot)$ be the distribution function of X + Y and let $J_{\ell}(\cdot)$ and $J^{u}(\cdot)$ be bounding functions such that $J_{\ell}(z) \leq J(z) \leq J^{u}(z)$ for all $z \in \mathbb{R}$.

Definition 7 (Achievability at a point). We say the lower bound $J_{\ell}(\cdot)$ is achievable at $z = z_0$ if there exists a joint distribution H of X, Y satisfying the marginals such that under H, $J(z_0) = J_{\ell}(z_0)$. The upper bound $J^u(\cdot)$ is achievable at $z = z_0$ if there exists a joint distribution H of X, Y satisfying the marginals such that under H, $J(z_0) = J^u(z_0)$.

Definition 8 (Pointwise Best-Possible). We say the lower bound $J_{\ell}(\cdot)$ is pointwise best-possible if for all $z_0 \in \mathbb{R}$ and $\varepsilon > 0$, $J_{\ell}(z_0) + \varepsilon$ will not be a valid lower bound for $J(z_0)$. In other words, for all $z_0 \in \mathbb{R}$ and $\varepsilon > 0$, there exists a joint distribution H of X, Y satisfying the marginals such that under H, $J(z_0) < J_{\ell}(z_0) + \varepsilon$. The upper bound $J^u(\cdot)$ is pointwise sharp if for all $z_0 \in \mathbb{R}$ and $\varepsilon > 0$, $J^u(z_0) - \varepsilon$ will not be a valid lower bound for $F(z_0)$.⁵

 $^{^{4}}$ Frank et al. (1987) shows that these bounds are equivalent to those given in Makarov (1982), which is the first paper to prove the bounds.

⁵Unlike Firpo and Ridder (2019), we differentiate between achievability and pointwise sharpness because a bound can be pointwise sharp but not necessarily achievable.

Definition 9 (Uniformly Sharp). We say the lower bound $J_{\ell}(\cdot)$ is uniformly sharp of H if there exists a single joint distribution H of X, Y satisfying the marginals such that under H, $J(z) = J_{\ell}(z)$ for all $z \in \mathbb{R}$. The upper bound $J^{u}(\cdot)$ is uniformly sharp if there exists a single joint distribution H of X, Y satisfying the marginals such that under H, $J(z) = J^{u}(z)$ for all $z \in \mathbb{R}$.

Following these definitions, if a bound is uniformly sharp, then it is achievable for all $z \in \mathbb{R}$ and also pointwise sharp. If a bound is achievable for all $z \in \mathbb{R}$, then it is pointwise sharp. However, a pointwise sharp bound may not be achievable for all $z \in \mathbb{R}$.

3.1 Prior results on achievability of bounds

We first state a theorem given in Nelsen (2006).⁶

Theorem 10 (Frank et al. Theorem 3.2, Nelsen Theorem 6.1.2). Let F and G be two fixed distribution functions. For any $z \in (-\infty, \infty)$:

(i) There exists a copula C_t , dependent only on the value t of $\tau_W(F,G)$ at z-, such that

$$\sigma_{C_t}(F,G)(z-) = \tau_W(F,G)(z-) = t$$
(13)

where z- is the left hand limit of the functions σ_{C_t} and τ_W as they approach z.⁷

(ii) There exists a copula C_r , dependent only on the value r of $\rho_W(F,G)(z)$, such that

$$\sigma_{C_r}(F,G)(z) = \rho_W(F,G)(z) = r.$$
(14)

Theorem 10 rephrases the result of Frank et al. (1987) in terms of the more common definition of right continuous distribution functions. The proof of Theorem 10 is sketched in Nelsen (2006). Embrechts et al. (2002) Theorem 5 also sketches the proof for the lower bound (13).

Kolmogorov's question is about the pointwise sharp bounds on the distribution functions of sum of two random variables. Statement (ii) of Theorem 10 along with Sklar's Theorem shows that the

⁶Frank et al. (1987) uses an example of a degenerate distribution to show that τ_W and ρ_W can be achieved for certain F, G (where the bounds are uniformly sharp in the example). However, it is not clear that for arbitrary F, G, Kolmogorov's question is answered.

⁷Frank et al. (1987) uses a left-continuous definition of the distribution functions. When translating the result, we need to be careful about the implications. In general, $\tau_W(\tilde{F}, \tilde{G})(z-) \leq \tau_W(\tilde{F}, \tilde{G})(z) = \tau_W(F, G)(z-) \leq \tau_W(F, G)(z)$.

Makarov upper bound (12) is achievable for all $z \in \mathbb{R}$ and thus is pointwise sharp. We will next consider two examples which show that statement (i) of Theorem 10 does not completely address Kolmogorov's question.

Example 11. Let X, Y be Bernoulli random variables with $p_1 = p_2 = 0.5$. Let F, G be the distribution functions of X, Y respectively. Suppose that we are interested in obtaining a lower bound for $P(X + Y \le 1)$. Clearly, $\tau_W(F,G)(1) = 0.5$ while $\tau_W(F,G)(1-) = 0$. Statement (i) of Theorem 10 does not provide bounds on $P(X + Y \le 1) = J(1)$.

Example 12 (Nelsen 2006). Let X, Y be random variables with uniform distributions on [0, 1]. Let F, G be distribution functions of X, Y respectively. Suppose that we wish to obtain a lower bound for $P(X + Y \le 1)$. In this example, $\tau_W(F, G)(1) = \tau_W(F, G)(1-) = 0$. We obtain the lower bound for $J(z) = P(X + Y \le 1) = 0$. Statement (i) of Theorem 10 tells us that there exists a joint distribution such that P(X + Y < 1) = 0. In fact, the construction in the proof of Theorem 10, given by Nelsen will result in $P(X + Y \le 1) = 1$ and P(X + Y < 1) = 0 when X = 1 - Y. However, this leaves open the question of whether $\tau_W(F, G)(1) = 0$ is a tight bound on $P(X + Y \le 1)$, either in the sense of being achievable, or more weakly, best possible.

To summarize, statement (i) of Theorem 10 leaves two questions unanswered: First, as shown in Example 11, when F, G are not continuous, $\tau_W(F, G)(z-)$ can be different from $\tau_W(F, G)(z)$. Thus when F and G are not continuous Theorem 10 (i) does not provide any information regarding whether the bound $\tau_W(F, G)(z)$ is either achievable or best possible vis a vis $J(z) = P(X + Y \leq z)$. Second, even when F, G are continuous, so that $\tau_W(F, G)(z-) = \tau_W(F, G)(z)$, it is still possible that $\sigma_{C_t}(F, G)(z-) < \sigma_{C_t}(F, G)(z)$. Consequently, even in the continuous case, the existence of the copula C_t given in statement (i) of Theorem 10 merely establishes that $\tau_W(F, G)(z-)$ is achievable as a lower bound on P(X + Y < z); it says nothing about the achievability (or otherwise) of $\tau_W(F, G)(z)$ as a lower bound on J(z).

3.2 When are the bounds achievable?

The next Theorem establishes that that for all F, G (possibly discontinuous) $\tau_W(F,G)(z)$ is best possible – thus addressing the unanswered question of Kolmogorov. We will return to the issue of achievability of $\tau_W(F,G)(z)$ as a lower bound on $P(X + Y \leq z)$, in Theorems 15 (discrete case) to 21 (general case) below.

Theorem 13 (No loose end to Kolmogorov's question). Let F and G be two fixed distribution functions. For any $z \in (-\infty, \infty)$, let $s = \tau_W(F, G)(z)$. For any $\varepsilon > 0$, there exists a copula $C_{s,\varepsilon}$ such that

$$\sigma_{C_{s,\varepsilon}}(F,G)(z) < s + \varepsilon.$$

In other words, the lower bound $\tau_W(F,G)(z)$ on J(z) cannot be improved for any F,G, and is thus pointwise best possible in terms of z.

Before proving Theorem 13, we first state a useful lemma.

Lemma 14 (Firpo and Ridder Theorem 2). For fixed distribution functions F, G, the function

$$\tau_W(F,G)(z) = \sup_{x+y=z} \max\{F(x) + G(y) - 1, 0\}$$

is right continuous and non-decreasing for all z.

Lemma 14 follows from the fact that F, G are right continuous and nondecreasing.

Proof. The proof of Theorem 13 uses a similar construction as in Makarov (1982). Now suppose that for a given z, $\tau_W(F,G)(z) = s$. Following Lemma 14, for all $\varepsilon > 0$, there exists $\delta > 0$ such that $\tau_W(F,G)(z+m) - s < \varepsilon$ for all $0 < m < \delta$. Pick any $0 < m < \delta$ that meets the above condition that $\tau_W(F,G)(z+m) < s + \varepsilon$. Then there exists a copula $C_{s,\varepsilon}$ such that

$$\sigma_{C_{s,\varepsilon}}(F,G)(z) \le \sigma_{C_{s,\varepsilon}}(F,G)((z+m)-) \tag{15}$$

$$= \tau_W(F,G)((z+m)-)$$
 (16)

$$\leq \tau_W(F,G)(z+m) \tag{17}$$

$$\langle s + \varepsilon$$
 (18)

where (15) follows from the fact that for any random variable D, $P(D \le d) \le P(D < d + h)$ for all h > 0; (16) follows from the optimality result in part (i) of Theorem 10; (17) follows from the non-decreasing property of τ_W ;(18) follows from the construction of $\tau_W(F,G)(z+m)$. This implies that for all $\varepsilon > 0$, we can construct a copula $C_{s,\varepsilon}$ such that $\sigma_{C_{s,\varepsilon}}(F,G)(z) < \tau_W(F,G)(z) + \varepsilon$, meaning that $\tau_W(F,G)(z) + \varepsilon$ will not be a valid lower bound on $P(X+Y \leq z) = J(z)$.

Theorem 13 along with Theorem 10 proves that Makarov's bounds are pointwise best-possible and thus directly address Komogorov's question. This closes the best-possible question left open by Theorem 10 (i).

3.3 Special case: at least one discrete variable

The following theorem states that the lower bound $\tau_W(F,G)(z)$ is achievable when we are dealing with one or more discrete random variables.

Theorem 15. If at least one of X, Y is a discrete random variable (a random variable which may take on only a countable number of distinct values), then there exists a copula C_t , dependent only on the value of $t = \tau_W(F, G)(z)$, such that

$$\sigma_{C_t}(F,G)(z) = \tau_W(F,G)(z) = t.$$

In other words, the lower bound on J(z) is always achievable when one of X, Y is a discrete random variable.⁸

Proof. Before proving the Theorem 15, we will first prove a useful lemma.

Lemma 16. For any k < z, if $t = \tau_W(F,G)(z) > \tau_W(F,G)(z-)$, there cannot be x',y' with x' + y' = k < z such that F(x') + G(y') - 1 = t.

Proof. Suppose that there exist x', y' with x' + y' = k < z and F(x') + G(y') - 1 = t, then by the

⁸Notice that this is Theorem 10 (i) with z- replaced by z, which can be surprising because Frank et al. (1987) note that theorem 10 (i) cannot be strengthened "even" when F, G are continuous, which implicitly implies that it cannot be strengthened when F, G are discrete. Some misconception about this point can be found in related literatures. For example, Kim (2014) stated that "If the marginal distributions of X and Y are both absolutely continuous with respect to the Lebesgue measure on \mathbb{R} , then the Makarov upper bound and lower bound can be achieved", which is contradicted by Example 12; Similarly, Williamson and Downs (1990) stated that in order for Theorem 10 to hold, it is necessary that F, G are not both discontinuous at a point x, y such that x + y = z; whereas Frank et al. (1987)'s proof does not require this to hold.



Figure 1: Support of C_t and mass assigned by C_t

definition of sup,

$$t \le \sup_{x+y=k} \max(F(x) + G(y) - 1, 0)$$
(19)

$$\leq \lim_{\varepsilon > 0, \varepsilon \to 0} \sup_{x+y=z-\varepsilon} \max(F(x) + G(y) - 1, 0)$$
(20)

$$=\tau_W(F,G)(z-) \tag{21}$$

$$\leq \sup_{x+y=z} \max(F(x) + G(y) - 1, 0) = t$$
(22)

where the inequalities (20) and (22) follow from the fact that the function $\tau_W(F,G)(\cdot)$ is nondecreasing and (19),(21) and (22) are by definition of $\tau_W(F,G)(k)$, $\tau_W(F,G)(z-)$, $\tau_W(F,G)(z)$. Thus, we have $\tau_W(F,G)(z) = \tau_W(F,G)(z-)$, which contradicts with the hypothesis that $\tau_W(F,G)(z) > \tau_W(F,G)(z-)$.

Now we will prove Theorem 15. Without loss of generality, we assume X is a discrete random variable. For $t = \tau_W(F, G)(z)$, we construct the copula

$$C_t(u,v) = \begin{cases} \operatorname{Max}(u+v-1,t), & (u,v) \text{ in } [t,1] \times [t,1], \\ \operatorname{Min}(u,v), & \text{otherwise.} \end{cases}$$

Figure 1 illustrates the support of C_t and the mass assigned by C_t .

Let $H_1 := \{(u, v) \in [0, 1]^2 \mid u + v - 1 = t\}, H_2 := \{(u, v) \in [0, 1]^2 \mid u = v < t\}, S_z := \{(u, v) \in [0, 1]^2 \mid F^{-1}(u) + G^{-1}(v) = z\}, S_{\bar{z}} := \{(u, v) \in [0, 1]^2 \mid F^{-1}(u) + G^{-1}(v) \le z\},$ where F^{-1}, G^{-1} are the

generalized inverses defined in Definition 2. Since t is a lower bound for $P(X + Y \leq z)$, C_t assigns mass at least t to the set $S_{\bar{z}}$. In particular, $H_2 \subseteq S_{\bar{z}}$. Thus, whether or not the lower bound on $P(X + Y \leq z)$ is achieved (by C_t) depends on whether the mass that C_t assigns to the set $H_1 \cap S_{\bar{z}}$ is 0.

We claim that $\iint_{S_{\overline{z}}\cap H_1} dC_t(u,v) = \iint_{S_z\cap H_1} dC_t(u,v)$. First suppose $\tau_W(F,G)(z) = \tau_W(F,G)(z-) = t$ t. Frank et al. (1987), showed that $P(X + Y < z) = \tau_W(F,G)(z-) = t$ under C_t . Thus, under C_t , the set $\{(u,v) \in [0,1]^2 | F^{-1}(u) + G^{-1}(v) < z\}$ will contain all the mass in H_2 (equals t) but not any mass in H_1 , so that the claim holds in this case. Now suppose $\tau_W(F,G)(z) > \tau_W(F,G)(z-)$, from Lemma 16 we know that the set $\{(u,v) \in [0,1]^2 | F^{-1}(u) + G^{-1}(v) < z\}$ cannot contain mass in H_1 . Therefore, $\iint_{S_{\overline{z}}\cap H_1} dC_t(u,v) = \iint_{S_z\cap H_1} dC_t(u,v)$, establishing the claim.

Since C_t assigns mass (1 - t) uniformly to H_1 , if $S_z \cap H_1$ is empty or only contains countably many points, then $\iint_{S_z \cap H_1} dC_t(u, v) = 0$, which is sufficient to establish the claim.

Since X is discrete, x can take at most countably many values with non-zero probability under F. For a given z, there are at most countably many points (x, y) such that x + y = z and P(X = x) > 0. Observe that

$$\{(u,v) \in [0,1]^2 | F^{-1}(u) + G^{-1}(v) = z\} = \bigcup_{(x,y): x+y=z} R_{xy}$$
(23)

where $R_{xy} \equiv \{(F(x-), F(x)] \times (G(y-), G(y))\}$ and we define the sets of form (a, a] as $\{a\}$ for any $a \in \mathbb{R}$, which will arise if Y is not discrete.

We will show that for each (x, y) with x + y = z, $R_{xy} \cap H_1$ contains at most one point. Since by definition of C_t , $t = \sup_{x+y=z} \max\{F(x) + G(y) - 1, 0\}$, we have $F(x) + G(y) - 1 \le t$ for any x+y=z. For any $(u,v) \in R_{xy}$ with u < F(x) or v < G(y), it holds that u+v-1 < t and thus (u,v)cannot be in H_1 . Therefore $R_{xy} \cap H_1$ contains at most one point (F(x), G(y)).⁹ As a consequence by (23), there exist at most countably many points in $S_z \cap H_1$. Thus, $\iint_{S_z \cap H_1} dC_t(u,v) = 0$ and

$$\sigma_{C_t}(F,G)(z) = \tau_W(F,G)(z) = t.$$

⁹Each rectangular region R_{xy} can touch the line u + v - 1 = t for at most one point because all points in R_{xy} satisfy $F^{-1}(u) + G^{-1}(v) = z$ and if there is more than one point in the intersection then t is not $\sup_{x+y=z} \max\{F(x) + G(y) - 1, 0\}$.

3.4 C_t not achieving the bound implies no other copula achieves the bound

Theorem 10 shows that the lower bound $\tau_W(F,G)(z-)$ on J(z-) can be achieved. In fact, the proof of Theorem 10 (see Frank et al. 1987 and Nelsen 2006) shows that the copula we constructed as C_t with $t = \tau_W(F,G)(z-)$ in the proof of Theorem 15 will achieve the lower bound $\tau_W(F,G)(z-)$.¹⁰ We further showed in Theorem 13 that the lower bound $\tau_W(F,G)(z)$ on J(z) cannot be improved. In example 12, for $t = \tau_W(F,G)(z) = \tau_W(F,G)(z-)$, we see that P(X + Y < z) = t under C_t but $P(X + Y \le z) > t$ under C_t .

This raises a new question: if we care not merely about sharpness, but also about the achievability of the lower bound $\tau_W(F,G)$ on J(z) – rather than J(z-) – and if C_t does not achieve the bound $\tau_W(F,G)$, can there be other copulas that can achieve the bound $\tau_W(F,G)$ for J(z)? Indeed, Frank et al. (1987) and Nelsen (2006) both pointed out that there are other copulas beside C_t that achieve the lower bound $\tau_W(F,G)(z-)$ for J(z-).

The corollary of the next theorem implies that for continuous F, G and an arbitrary z, in order to determine whether the lower bound $\tau_W(F, G)(z)$ on J(z) can be achieved, we only need to determine whether it is achieved under C_t for $t = \tau_W(F, G)(z)$. Theorem 19 along with Theorem 17 further establishes this claim for arbitrary F, G. In other words, if the lower bound for J(z) is not achieved under C_t , then there is no joint distribution that will achieve this lower bound.

Theorem 17. Given arbitrary z and F, G, if $\tau_W(F, G)(z-) = \tau_W(F, G)(z) = t$ and the copula C_t does not achieve the lower bound $\tau_W(F, G)(z)$ on $J(z) \equiv P(X + Y \leq z)$, then no other copula can achieve this lower bound.

Proof. Since $\tau_W(F,G)(z-) = t$, by Theorem 10(i), copula C_t achieves the bound $\tau_W(F,G)(z-)$ on P(X + Y < z). That is, C_t assigns mass t to the set $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) < z\}$. Since, by hypothesis, C_t does not achieve the lower bound $\tau_W(F,G)(z)$ on $P(X + Y \le z)$, C_t assigns non-zero probability to the set $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) = z\}$. In particular, the image of the set $\{(x, y) : x + y = z\}$ under the (F, G) mapping must contain a line segment with

¹⁰In fact, as noted above, Frank et al. (1987) consider bounds on \tilde{J} , Nelsen (2006) translate the result to the standard definition J but do not provide a full proof.

length greater than 0 on the line u + v - 1 = t in the uv-plane inside the unit square¹¹ as illustrated in Figure 2; since otherwise under C_t , $P(X + Y < z) = P(X + Y \le z)$ in which case C_t achieves the bound. Let a, b be such that the line segment $\{(u, v) : u = a + s, v = 1 - t - (a + s) \text{ for } s, 0 \le s \le b - a\}$ is contained in $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) = z\} \cap \{(u, v) \subseteq [0, 1] \times [0, 1] : u + v - 1 = t\}$.

Now suppose there is a copula C that achieves the lower bound $\tau_W(F,G)(z)$ on $P(X + Y \leq z)$. First, we claim that C must assign mass t to the rectangle $R_1 = [0, a] \times [0, 1 + t - b]$. Since R_1 is a subset of $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) \leq z\}$ and by hypothesis C achieves the lower bound, C cannot assign mass more than t to R_1 .

Suppose C assigns mass 0 < r < t to R_1 ; see Figure 3. Note that we define the margins of the copula to be uniform (C(p, 1) = C(1, p) = p, for all p in [0, 1]). In particular, in order for C(1, 1 + t - b) = 1 + t - b, C needs to assign mass 1 + t - b - r to $[a, 1] \times [0, 1 + t - b]$ and similarly C needs to assign mass a - r to $[0, a] \times [1 + t - b, 1]$. As a consequence, C needs to assign mass 1 - (1 + t - b - r) - (a - r) - r = b - a + r - t to the rectangle $[a, 1] \times [1 + t - b, 1]$. Now consider the rectangle $[a, b] \times [0, 1 + t - b]$. It needs to contain mass at least t - r since $[a, b] \times [0, 1]$ needs to contain mass b - a and $[a, b] \times [1 + t - b, 1] \subseteq [a, 1] \times [1 + t - b, 1]$. Similarly, $[0, a] \times [1 + t - b, 1 + t - a]$ needs to contain mass at least t - r. Then C assigns mass greater than or equal to r + 2(t - r) = t + (t - r) > tto $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) \le z\}$, which is a contradiction that C achieves the lower bound $\tau_W(F, G)(z)$ on $P(X + Y \le z)$. Therefore, C must assign mass t to the rectangle $R_1 = [0, a] \times [0, 1 + t - b]$.

Next, we show that C assigns mass b - a to the rectangle $[a, b] \times [1 + t - b, 1 + t - a]$. By the hypothesis that C achieves the lower bound $\tau_W(F,G)(z)$ on $P(X + Y \leq z)$, C assigns mass 1 - t to the set $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) > z)\}$, which is a subset of the union of the following three rectangles: $[0, 1] \times [1 + t - a, 1], [a, b] \times [1 + t - b, 1 + t - a], [b, 1] \times [0, 1]$. In order to maintain uniform margins, the first and third rectangles contain mass a - t and 1 - b. So the rectangle $[a, b] \times [1 + t - b, 1 + t - a]$ needs to contain mass at least (1 - t) - (a - t) - (1 - b) = b - a. Again from the uniformity of the margins, $[a, b] \times [1 + t - b, 1 + t - a]$ can contain mass at most b - a. Thus, C assigns mass b - a to the rectangle $[a, b] \times [1 + t - b, 1 + t - a]$. Furthermore, since the rectangle $[a, 1] \times [1 + t - b, 1]$ contains total mass 1 - (1 + t - b) - (a - t) = b - a, there's no mass

¹¹The first two sentences of the proof imply that when $\tau_W(F,G)(z) = \tau_W(F,G)(z-)$ this is a necessary and sufficient condition for C_t to assign non-zero probability to the set $\{(u,v) \subseteq [0,1] \times [0,1] : F^{-1}(u) + G^{-1}(v) = z\}$.

elsewhere in this rectangle except in $[a, b] \times [1 + t - b, 1 + t - a]$.

Now we show that C needs to assign mass b-a to the line segment (a, 1+t-a) to (b, 1+t-b)inside the square $[a, b] \times [1 + t - b, 1 + t - a]$. Figure 3 shows a zoomed-in version of the rectangle $[a,b] \times [1+t-b,1+t-a]$. First, C cannot assign any mass strictly below the line segment (a,1+t-a)to (b, 1+t-b) inside the square $[a, b] \times [1+t-b, 1+t-a]$ because C already assigns mass t to R_1 and the total mass assigned by C_t to the set $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) < z\}$ is t. For any $\text{rectangle } [m,n] \times [c,d] \text{ such that } m+c-1 \geq t, \, a \leq m < n \leq b, 1+t-b \leq c < d \leq 1+t-a, \, \text{suppose } a \leq t, t \in t, t \int, t, t \int$ that C assigns mass $\delta > 0$ to this rectangle. We know that C assigns mass t to the region R_1 . Let E_1 be the triangular area defined by vertices (m, 1+t-m), (a, 1+t-a), (m, 1+t-a) and E_2 be the triangular area defined by vertices (m, 1+t-m), (b, 1+t-b), (b, 1+t-m), as depicted in Figure 3. Note that we have previously established that C assigns no mass to the rectangle $[a, b] \times [1 + t - a, 1]$. The mass assigned by C to E_1 and R_1 is equal to the mass in rectangle $[0, m] \times [0, 1]$ subtracting the mass in the rectangle $[0, a] \times [1+t-a, 1]$, which is C(m, 1) - (a-t) = m - (a-t) = m - a + t. Similarly, the mass assigned by C to E_2 and R_1 is equal to C(1, 1+t-m)-(1-b) = 1+t-m-(1-b) = t-m+b. Thus, C assigns the mass m-a to E_1 and b-m to E_2 . Since E_1, E_2 are disjoint, C assigns the mass b-a to $E_1 \cup E_2$. Then the rectangle $[m, n] \times [c, d]$ will contain mass 0, which is a contradiction. Since the choice of $[m, n] \times [c, d]$ is arbitrary, we know that C assigns mass b - a to the line segment in \mathbb{R}^2 from (a, 1-t-a) to (b, 1-t-b) inside the square $[a, b] \times [1+t-b, 1+t-a]$. Finally, C assigns mass at least t + (b-a) to the set $\{(u, v) \subseteq [0, 1] \times [0, 1] : F^{-1}(u) + G^{-1}(v) \leq z\}$, which contradicts that C achieves the lower bound $\tau_W(F,G)(z)$ on $P(X+Y \leq z)$. Thus, when $\tau_W(F,G)(z) = \tau_W(F,G)(z-)$, if the copula C_t does not achieve the lower bound $\tau_W(F,G)(z)$ on $P(X+Y \leq z)$, then no other copula can achieve this lower bound.

Corollary 18. Given arbitrary z and continuous F, G, let $t = \tau_W(F, G)(z)$. If the copula C_t does not achieve the lower bound $\tau_W(F, G)(z)$ on $J(z) \equiv P(X + Y \leq z)$, then no other copula can achieve this lower bound.

Proof. This follows directly from Theorem 17 since when F, G are both continuous, $\tau_W(F, G)(z) = \tau_W(F, G)(z-)$.



Figure 2: Copula C and the image of x+y = zunder the (F, G) mapping.



Figure 3: A zoom in part of Figure 2 with rectangle $[m, n] \times [c, d]$ colored in blue.

3.5 Sufficient conditions for achievability of the lower bound on J(z)

Theorem 19. Given arbitrary z and F, G, if $\tau_W(F, G)(z) > \tau_W(F, G)(z-)$ then the copula C_t with $t = \tau_W(F, G)(z)$ will achieve the lower bound $\tau_W(F, G)(z)$ on $P(X + Y \leq z)$.

Proof. We will prove the contrapositive: if the lower bound $t = \tau_W(F,G)(z)$ of $P(X+Y \leq z)$ is not achievable under C_t , then $\tau_W(F,G)(z) = \tau_W(F,G)(z-)$. Note that $\tau_W(F,G)(z) = \tau_W(F,G)(z-)$ holds trivially when $\tau_W(F,G)(z) = 0$. We will assume $t = \tau_W(F,G)(z) > 0$.

If the lower bound $\tau_W(F,G)(z)$ of $P(X + Y \leq z)$ is not achievable under C_t , the set $\{(u,v) \in [0,1]^2 | F^{-1}(u) + G^{-1}(v) \leq z\}$ must contain a line segment with length greater than 0 on the line u + v - 1 = t in the *uv*-plane. Based on Lemma 16, the set $\{(u,v) \in [0,1]^2 | F^{-1}(u) + G^{-1}(v) = z\}$ must contain a line segment with length greater than 0 on the line u + v - 1 = t in the *uv*-plane. The existence of this line segment implies that the image of the set $\{(x,y) : x + y = z\}$ under the (F,G) mapping must also contain a line segment with length greater than 0 on the line u + v - 1 = t in the u + v - 1 = t in the *uv*-plane. This means that there exist x^* and $\varepsilon > 0$ such that F(x) + G(z - x) - 1 = t for all $x \in (x^* - \varepsilon, x^* + \varepsilon)$ and $F(\cdot)$ is continuous and strictly increasing on $x \in (x^* - \varepsilon, x^* + \varepsilon)$. In particular, for any $\delta > 0$, there exists $\varepsilon^* > 0$ such that $F(x^*) - F(x^* - \varepsilon^*) < \delta$. By definition of x^* , $\tau_W(F,G)(z) = F(x^*) + G(z - x^*) - 1$. Then for $\varepsilon^* > 0$,

$$\tau_W(F,G)(z-\varepsilon^*) = \sup_{x+y=z-\varepsilon^*} \max(F(x) + G(y) - 1, 0) \ge F(x^* - \varepsilon^*) + G(z-x^*) - 1$$
(24)

And

$$\tau_W(F,G)(z) - \tau_W(F,G)(z - \varepsilon^*) = F(x^*) + G(z - x^*) - 1 - \tau_W(F,G)(z - \varepsilon^*)$$
(25)

$$\leq F(x^*) + G(z - x^*) - 1 - (F(x^* - \varepsilon^*) + G(z - x^*) - 1)$$
 (26)

$$=F(x^*) - F(x^* - \varepsilon^*) < \delta \tag{27}$$

Since δ is arbitrary, $\tau_W(F, G)(\cdot)$ is continuous at z and we must have $\tau_W(F, G)(z) = \tau_W(F, G)(z-)$.

Thus it follows from Theorem 19 that the only time when the lower bound on $P(X + Y \le z)$ is not achievable is when the pointwise best possible bounds for $P(X + Y \le z)$ and P(X + Y < z)are the same. This result can be quite surprising: it follows that the distribution implied for X + Yvia the construction of C_t , namely, $\sigma_{C_t}(F,G)(z) = P(X + Y \le z)$, is discontinuous at z only when $\tau_W(F,G)(z)$ is continuous at z, i.e. $\tau_W(F,G)(z-) = \tau_W(F,G)(z)$.

We present an example to show that we do not require the margins F, G to be cdf of uniform distributions for the lower bound $\tau_W(F, G)(z)$ on $P(X + Y \leq z)$ to be not achievable.

Example 20. Let

$$F(x) = \begin{cases} 0 & x < 0 \\ x^2 & 0 \le x < 1 \\ 1 & x \ge 1 \end{cases} \qquad G(y) = \begin{cases} 0 & y < 0 \\ 1 - (1 - y)^2 & 0 \le y < 1 \\ 1 & y \ge 1 \end{cases}$$

F is the distribution for random variable X where X follows a triangular distribution with a = 0, b = c = 1 (equivalent to Beta(2,1)) and G is the distribution for random variable Y where Y follows a triangular distribution with a = c = 0, b = 1 (equivalent to Beta(1,2)). Suppose z = 1. Then

$$\tau_W(F,G)(1) = \sup_{x+y=1} \max\{F(x) + G(y) - 1, 0\}$$
(28)

$$= \sup_{x} \max\{F(x) + G(1-x) - 1, 0\}$$
(29)

$$=0$$
(30)

The lower bound equals 0 corresponds to the copula C_0 constructed based on the lower Fréchet-Hoeffding bound (in other words, X, Y are perfectly negatively correlated). In this example, X = 1 - Y. So under C_0 , $P(X+Y \le 1) = 1$ and P(X+Y < 1) = 0. The lower bound $\tau_W(F,G)(1)$ on $P(X+Y \le 1)$ is not achievable.

3.6 Characterization of achievability of the lower bound on J(z)

Theorem 21. The Makarov lower bound $t = \tau_W(F,G)(z)$ on $P(X + Y \le z)$ is not achievable at zif and only if there exists x^*, y^* with $x^* + y^* = z$ such that all following three conditions hold: (i) $F(x^*) + G(y^*) = \sup_{x+y=z} \{F(x) + G(y)\} \ge 1$; (ii) F(x) + G(y) is constant for x in a neighborhood $Nr(x^*)$ of x^* and y = z - x; (iii) the image of the set $\{x, y : x \in Nr(x^*), y = z - x\}$ under the (F, G)mapping contains an open interval within the line segment $\{(u, v) \in [0, 1]^2 \mid u + v - 1 = t\}$.¹²

Proof. If the Makarov lower bound $\tau_W(F,G)(z) =: t$ is not achievable at z, then it is not achievable under C_t . In particular, for $S_z := \{(u,v) \in [0,1]^2 | F^{-1}(u) + G^{-1}(v) = z\}$ and $H_1 := \{(u,v) \in [0,1]^2 | u+v-1=t\}$, $P(X+Y=z) = \iint_{S_z \cap H_1} dC_t(u,v) > 0$.

If (i) does not hold, then there are two sub-cases: if $\sup_{x+y=z} \{F(x) + G(y)\} < 1$, then $t = \tau_W(F,G)(z) = 0$, hence the set S_z does not intersect H_1 and thus $S_z \cap H_1 = \emptyset$. On the other hand, if there do not exist x^*, y^* such that $F(x^*) + G(y^*) = \sup_{x+y=z} \{F(x) + G(y)\}$, then $S_z \cap H_1 = \emptyset$ by definition. In both sub-cases, the lower bound is achieved under C_t , which is a contradiction.

Now suppose that (i) holds. Since C_t assigns mass uniformly to the set H_1 , and by hypothesis, $\iint_{S_z \cap H_1} dC_t(u, v) > 0$, there exists a neighborhood of $(F(x^*), G(y^*))$ in H_1 that is contained in S_z . Further, this neighborhood of $(F(x^*), G(y^*))$ in H_1 is contained in the image of $\{(x, y) \mid x + y = z\}$ under the (F, G) mapping. Thus, (iii) holds and (ii) holds by definition of H_1 .

For the converse, if (i), (ii), (iii) hold, then S_z contains a non-zero measure set in H_1 . Since C_t assigns mass uniformly to the set $H_1, \iint_{S_z \cap H_1} dC_t(u, v) > 0$ and the lower bound $\tau_W(F, G)(z)$ will not be achievable at z under C_t . Furthermore, by Theorem 19, (i), (ii), (iii) imply that $\tau_W(F, G)(z) =$ $\tau_W(F, G)(z-)$. Finally by Theorem 17, the lower bound $\tau_W(F, G)(z)$ on $P(X + Y \leq z)$ will not be achievable for any copula.

¹²The conditions can also be defined similarly using the neighborhood of y^* .

Theorem 22. Makarov bounds on $P(X + Y \le z)$ are pointwise best-possible, ¹³ but the lower bound may not be achievable for all $z \in \mathbb{R}$. The Makarov upper bound is achievable for all $z \in \mathbb{R}$. Makarov bounds are in general not uniformly sharp¹⁴.

The following table summarizes when the Makarov bounds are always achievable (for all $z \in \mathbb{R}$ and any F, G) under different definitions of distribution functions.

	P(X + Y < z)	$P(X+Y \le z)$
Upper Bound		\checkmark
Lower Bound	\checkmark	

Table 1: The Makarov upper bound on P(X+Y < z) and the Makarov lower bound on $P(X+Y \le z)$ are always achievable for any given marginals F, G and for all $z \in \mathbb{R}$. The achievabilities of the Makarov upper bound on P(X+Y < z) and the Makarov lower bound on $P(X+Y \le z)$ are margin specific and depend on z. See Example 12 and Theorem 21.

4 Sharp bounds on the difference

Now we consider the case of the difference of two random variables with fixed marginals. We will make the substitution when one variable is negated to get the best-possible bounds on the difference. Let X and Y be random variables with respective distribution functions F and G fixed. Let $\Delta = X - Y$ be the difference of random variables X, Y. Let $J_{\Delta}(\cdot)$ be the distribution function of Δ .

Theorem 23. For any given value δ , best-possible bounds on $J_{\Delta}(\delta)$ where $\underline{J_{\Delta}}(\delta) \leq J_{\Delta}(\delta) \leq \overline{J_{\Delta}}(\delta)$ are given by

$$\underline{J}_{\underline{\Delta}}(\delta) = \sup_{x-y=\delta} \max\{F(x) - P(Y < y), 0\}$$
$$= \sup_{x-y=\delta} \max\{F(x) - G(y) + P(Y = y), 0\}$$
(31)

$$\overline{J_{\Delta}}(\delta) = 1 + \inf_{x-y=\delta} \min\{F(x) - P(Y < y), 0\}$$

= $1 + \inf_{x-y=\delta} \min\{F(x) - G(y) + P(Y = y), 0\}$ (32)

 $^{13}\mathrm{See}$ Definition 8

¹⁴For given F, G, there does not exist a single joint distribution that achieves the bounds for all z. For example, the construction of the copula C_t depends on the value z in general.

The bounds in Theorem 23 differ from Fan and Park (2010) and Williamson and Downs (1990) where we have the point mass P(Y = y) in both the upper and lower bounds.

4.1 Proof of Theorem 23

Consider a new variable Y' = -Y with cdf G'. Then from equation (8) and (12), for any δ , the bound on $P(\Delta \leq \delta) = P(X - Y \leq \delta) = P(X + Y' \leq \delta)$ is:

$$\underline{J_{\Delta}}(\delta) = \sup_{x+y'=\delta} \max(F(x) + G'(y') - 1, 0)$$
(33)

$$\overline{J_{\Delta}}(\delta) = 1 + \inf_{x+y'=\delta} \min(F(x) + G'(y') - 1, 0)$$
(34)

Note that

$$G'(y') = P(-Y \le y') \tag{35}$$

$$=P(Y \ge -y') \tag{36}$$

$$= 1 - P(Y < -y') \tag{37}$$

$$= 1 - G(-y') + P(Y = -y')$$
(38)

Replace y' with -y and G'(y') with 1-G(y)+P(Y=y), we get the best-possible bounds in Theorem 23.

Remark 24. In the case where P(Y = y) = 0 for all y (for example, when the distribution function of Y is absolutely continuous with respect to the Lebesgue measure), we recover the bestpossible bounds in Theorem 2 of Williamson and Downs (1990) and in Lemma 2.1 of Fan and Park (2010). However, when G is not absolutely continuous, the bounds in Fan and Park (2010) and Williamson and Downs (1990) can be different from the bounds in Theorem 23 as the point mass P(Y = y) can be nonzero for some $y \in \mathbb{R}$.

4.2 Implications of the new bounds

What went wrong? In the proof of Theorem 2 of Williamson and Downs (1990), they stated that "Let Y' = -Y. Then G'(y) = 1 - G(-y)". This statement is not correct as in our notation,

$$G'(y) = P(Y' \le y) \tag{39}$$

$$= P(-Y \le y) \tag{40}$$

$$=P(Y \ge -y) \tag{41}$$

$$= 1 - P(Y < -y)$$
 (42)

$$= 1 - G(-y) + P(Y = -y)$$
(43)

If we use the left-continuous version definition of cdf in Williamson and Downs (1990) where $\tilde{G}'(y) = P(Y' < y)$, we also get

$$\tilde{G}'(y) = P(Y' < y) \tag{44}$$

$$= P(-Y < y) \tag{45}$$

$$=P(Y>-y) \tag{46}$$

$$= 1 - P(Y \le -y) \tag{47}$$

$$= 1 - \tilde{G}(-y) - P(Y = -y)$$
(48)

In fact, it is easy to show that G'(y) = 1 - G(-y) if and only if $\tilde{G}'(y) = 1 - \tilde{G}(-y)$. Although we only focus on the sum and difference of two random variables in this paper, the same mistake appears in the division of two random variables in Williamson and Downs (1990). At this point, Theorem 23 seems to imply that the lower bound in Williamson and Downs (1990) is valid but not necessarily best-possible and the upper bound might not be valid. The next theorem will establish the validity of the upper bound in Williamson and Downs (1990) even though the proof used in Williamson and Downs (1990) is not correct. **Theorem 25.** For any random variables X and Y with respective cdfs $F(\cdot)$ and $G(\cdot)$,

$$\inf_{x-y=\delta} \min\{F(x) - P(Y < y), 0\} = \inf_{x-y=\delta} \min\{F(x) - G(y), 0\}$$
(49)

Theorem 25 implies that the upper bounds in Williamson and Downs (1990) and Fan and Park (2010) coincides with the bounds we proposed in Theorem 23. Since the lower bounds that we propose in Theorem 23 is greater than or equal to the lower bounds in Williamson and Downs (1990), Fan and Park (2010), Theorem 25 establishes the validity of all these bounds. However, these lower bounds may not be optimal.

4.3 **Proof of Theorem** 25

We first prove some propositions that are used to prove Theorem 25. Let $f(\cdot)$ be a function that is non-decreasing and is right continuous, so that for all x:

$$\lim_{\varepsilon \to 0, \varepsilon > 0} f(x + \varepsilon) = f(x)$$
(50)

Let $g(\cdot)$ be a function that is non-increasing and is left continuous, so that for all x:

$$\lim_{\varepsilon \to 0, \varepsilon > 0} g(x - \varepsilon) = g(x) \tag{51}$$

Now consider the function $h(x) \equiv f(x) + g(x)$.

Proposition 26. For all x, we have:

$$h(x) \geq \lim_{\varepsilon \to 0, \varepsilon > 0} h(x + \varepsilon),$$
 (52)

$$h(x) \geq \lim_{\varepsilon \to 0, \varepsilon > 0} h(x - \varepsilon).$$
 (53)

Proof.

$$\lim_{\varepsilon \to 0, \varepsilon > 0} h(x + \varepsilon) = \lim_{\varepsilon \to 0, \varepsilon > 0} \left(f(x + \varepsilon) + g(x + \varepsilon) \right)$$
(54)

$$= f(x) + \lim_{\varepsilon \to 0, \varepsilon > 0} g(x + \varepsilon)$$
(55)

$$\leq f(x) + g(x) = h(x). \tag{56}$$

Here the second equality follows by right continuity of $f(\cdot)$ and the inequality follows since $g(\cdot)$ is non-increasing. Similarly:

$$\lim_{\varepsilon \to 0, \varepsilon > 0} h(x - \varepsilon) = \lim_{\varepsilon \to 0, \varepsilon > 0} \left(f(x - \varepsilon) + g(x - \varepsilon) \right)$$
(57)

$$= \lim_{\varepsilon \to 0, \varepsilon > 0} f(x - \varepsilon) + g(x)$$
(58)

$$\leq f(x) + g(x) = h(x). \tag{59}$$

Here the second equality is by left continuity of $g(\cdot)$ and the inequality follows since $f(\cdot)$ is nondecreasing.

Now consider the right continuous non-increasing function g^* defined as follows:

$$g^{*}(x) = \begin{cases} g(x) & \text{if } g(x) = \lim_{\varepsilon \to 0, \varepsilon > 0} g(x + \varepsilon) \\ \lim_{\varepsilon \to 0, \varepsilon > 0} g(x + \varepsilon) & \text{if } g(x) > \lim_{\varepsilon \to 0, \varepsilon > 0} g(x + \varepsilon) \end{cases}$$
(60)

In words, if x is a point of discontinuity of $g(\cdot)$ then $g^*(x)$ is defined to be the right hand limit of $g(\cdot)$, while $g(x) = g^*(x)$ at points of continuity of $g(\cdot)$.

Proposition 27. For all x:

$$\lim_{\varepsilon \to 0, \varepsilon > 0} g(x + \varepsilon) = \lim_{\varepsilon \to 0, \varepsilon > 0} g^*(x + \varepsilon)$$
(61)

$$\lim_{\varepsilon \to 0, \varepsilon > 0} g(x - \varepsilon) = \lim_{\varepsilon \to 0, \varepsilon > 0} g^*(x - \varepsilon).$$
(62)

Proof. This follows from the definition of $g^*(\cdot)$.

Let $h^{*}(x) \equiv f(x) + g^{*}(x)$.

Proposition 28.

$$\inf_{y} h(y) = \inf_{y} h^{*}(y).$$
(63)

Proof: This follows from Proposition 26 and Proposition 27.

4.4 Proof of Theorem 25

Theorem 25 follows from Proposition 28 where $F(x) = Pr(X \le x)$ is right continuous and nondecreasing, -Pr(Y < y) is left continuous and non-increasing. Thus the left hand side of (49) corresponds to min{ $\inf_y h(y), 1$ }, while the right hand side of (49) corresponds to min{ $\inf_y h^*(y), 1$ }.

Remark 29. Theorem 3 Williamson and Downs (1990) which states an optimality result (analogous to Theorem 10 of our paper) contains an unnecessary exclusion: specifically, it states that the bounds are only achievable if F and G are not both discontinuous at some x, y such that x + y = z. It appears that this additional unnecessary condition was added because Williamson and Downs fail to note that when F and G are discontinuous the bounds only take a strict subset of values in [0,1]. Consequently, only a subset of values need to be considered.

5 A causal perspective

Throughout this section, we consider a binary treatment D = 0, 1. Let Y_1 be the potential outcome of receiving the treatment and Y_0 be the potential outcome of not receiving the treatment. We assume the stable unit treatment value assumption (SUTVA, Rubin 1978) that there is a single version of each treatment/control and no interference among the subjects. We define our parameter of interest $\Delta = Y_1 - Y_0$ (also regarded as the individual treatment effect). Fan and Park (2010) Lemma 2.1 stated sharp bounds on the distribution function of the individual treatment effect. We modify the bounds in Fan and Park (2010) based on Theorem 23. Let F_1, F_0 be the cumulative distribution function on Y_1, Y_0 respectively. Let $F_{\Delta}(\cdot)$ be the cdf for Δ . **Theorem 30.** For any given value δ , best-possible bounds on $F_{\Delta}(\delta)$ are given by

$$F^{L}(\delta) = \sup_{y} \max\{F_{1}(y) - P(Y_{0} < y - \delta), 0\}$$

=
$$\sup_{y} \max\{F_{1}(y) - F_{0}(y - \delta) + P(Y_{0} = y - \delta), 0\}$$
 (64)

$$F^{U}(\delta) = 1 + \inf_{y} \min\{F_{1}(y) - P(Y_{0} < y - \delta), 0\}$$

= 1 + \inf \min_{y} \min_{1}\{F_{1}(y) - F_{0}(y - \delta) + P(Y_{0} = y - \delta), 0\} (65)

Let Y denotes the observed variable. Under consistency, $Y = Y_0$ when D = 0 and $Y = Y_1$ when D = 1. In practice, if we are willing to assume ignorability or conditional ignorability (for example, in randomized clinical trials (RCTs)), the marginal distributions $F_1(y)$ and $F_0(y)$ can be identified. Theorem 30 allows us to conclude best-possible bounds on the distribution function of the individual treatment effect. In the special case where Y is ordinal, Proposition 1 in Lu et al. (2018) can be recovered using Theorem 23 and Theorem 25. Lu et al. (2018) considers a special case where Y is non-negative and proves the bounds using a construction argument instead of the copula theory.

Corollary 31. The Fan-Park upper bound is best-possible.

Proof: This follows directly from Theorem 25 where X is replace by Y_1 and Y is replace by Y_0 . \Box

5.1 Application of Theorem 30 on cdf bounds of ITE

Here we will present a simple example that applies our bounds in Theorem 30 and compare it with the bounds in Fan and Park (2010).

Consider the case where we have a binary treatment variable (D = 0, 1) and a ternary response (Y = 0, 1, 2). Under randomization, the relationship between the counterfactual distribution $P(Y_0, Y_1)$ and the observed distributions $\{P(Y \mid D = 0), P(Y \mid D = 1)\}$ is given by: $P(Y = i \mid D = j) = P(Y_j = i)$. Suppose we observe the marginals given in Table 2. We can parameterize the joint distribution with 4 parameters p, q, t, r.

	$P(Y\!=\!0 \mid D\!=\!0) = 0.3$	$P(Y\!=\!1 \mid D\!=\!0) = 0.2$	$P(Y\!=\!2 \mid D\!=\!0) = 0.5$
$P(Y=0 \mid D=1) = 0.7$	$P(Y = 0 \mid D = 1) - p - r$	$p \in [0, 0.2]$	$r \in [0.2, 0.5]$
$P(Y\!=\!1 \mid D\!=\!1) = 0.1$	$P(Y = 1 \mid D = 1) - t - q$	$t \in [0, 0.1]$	$q \in [0, 0.1]$
$P(Y\!=\!2 \mid D\!=\!1) = 0.2$	1 - ()	$P(Y = 1 \mid D = 0) - t - p$	$P(Y = 2 \mid D = 0) - r - q$

Table 2: Application with binary treatment and ternary outcome

Based on the bounds proposed in (64) and (65), we note the following alternative expressions for $F^{L}(\delta)$ and $F^{U}(\delta)$:

$$F^{L}(\delta) = \max\left(\sup_{y} \{F_{1}(y) - P(Y_{0} < y - \delta)\}, 0\right)$$
$$F^{U}(\delta) = 1 + \min\left(\inf_{y} \{F_{1}(y) - P(Y_{0} < y - \delta)\}, 0\right)$$

Consider $\delta = -2$, only possible when $Y_1 = 0, Y_0 = 2$. So this corresponds to the entry top right corner of Table 2. By Fréchet inequalities, the bounds on $P(Y_1 = 0, Y_0 = 3)$ is given by $r \in [0.2, 0.5]$. Now consider $F_1(y) - P(Y_0 < y - \delta)$ in our example,

$$F_{1}(y) - P(Y_{0} < y + 2) = \begin{cases} 0 & y \leq -2 \\ -0.3 & -2 < y \leq -1 \\ -0.5 & -1 < y < 0 \\ 0.2 & y = 0 \\ -0.3 & 0 < y < 1 \\ -0.2 & 1 \leq y < 2 \\ 0 & y \geq 2 \end{cases}$$

This gives bound for $F_{\Delta}(-2) \in [0.2, 0.5]$, which matches the Fréchet inequality bound. In this case

if we consider the bounds proposed in Lemma 2.1 in Fan and Park (2010),

$$F_1(y) - F_0(y+2) = \begin{cases} 0 & y < -2 \\ -0.3 & -2 \le y < -1 \\ -0.5 & -1 \le y < 0 \\ -0.3 & 0 \le y < 1 \\ -0.2 & 1 \le y < 2 \\ 0 & y \ge 2 \end{cases}$$

The lower bounds for $F_{\Delta}(-2)$ is 0, which is not sharp. This example corresponds to the case that F_1 and F_0 are both discontinuous at $Y_1 = 0$ and $Y_0 = 2$.

To complete this example, we calculate the bounds in Theorem 30 for $\delta = -1$, this corresponds to p + q + r. Now consider $F_1(y) - P(Y_0 < y + 1)$ in our example,

$$F_1(y) - P(Y_0 < y + 1) = \begin{cases} 0 & y \le -1 \\ -0.3 & -1 < y < 0 \\ 0.4 & y = 0 \\ 0.2 & 0 < y < 1 \\ 0.3 & y = 1 \\ -0.2 & 1 < y < 2 \\ 0 & y \ge 2 \end{cases}$$

This gives bounds for $F_{\Delta}(-1) = p + q + r \in [0.4, 0.7]$. Similarly, we can obtain the bounds for $F_{\Delta}(0) = 1.2 + t - r \in [0.7, 1]$, $F_{\Delta}(1) = 1.5 - p - q - t - r \in [0.8, 1]$. $F_{\Delta}(2) = 1$ follows trivially by construction. Kreinovich and Ferson (2006) discussed some computational perspectives of calculating the type of bounds in Theorem 23. Inference on Makarov type of bounds can be found in Fan and Park (2010) and is discussed in Fan and Park (2012) and Imbens and Menzel (2018).

6 Acknowledgments

We thank Carlos Cinelli, Yanqin Fan and audience from UW causal reading group / poster session for the valuable input and discussion.

References

- Chiba, Y. (2017). Sharp nonparametric bounds and randomization inference for treatment effects on an ordinal outcome. *Statistics in medicine*, 36(25):3966–3975.
- Embrechts, P. and Hofert, M. (2013). A note on generalized inverses. Mathematical Methods of Operations Research, 77(3):423–432.
- Embrechts, P., McNeil, A., and Straumann, D. (2002). Correlation and dependence in risk management: properties and pitfalls. *Risk management: value at risk and beyond*, 1:176–223.
- Fan, Y. and Park, S. S. (2010). Sharp bounds on the distribution of treatment effects and their statistical inference. *Econometric Theory*, 26(3):931–951.
- Fan, Y. and Park, S. S. (2012). Confidence intervals for the quantile of treatment effects in randomized experiments. *Journal of Econometrics*, 167(2):330–344.
- Firpo, S. and Ridder, G. (2019). Partial identification of the treatment effect distribution and its functionals. *Journal of Econometrics*, 213(1):210–234.
- Frank, M. J., Nelsen, R. B., and Schweizer, B. (1987). Best-possible bounds for the distribution of a sum—a problem of kolmogorov. *Probability theory and related fields*, 74(2):199–211.
- Huang, E. J., Fang, E. X., Hanley, D. F., and Rosenblum, M. (2017). Inequality in treatment benefits: Can we determine if a new treatment benefits the many or the few? *Biostatistics*, 18(2):308–324.
- Imbens, G. and Menzel, K. (2018). A causal bootstrap. Technical report, National Bureau of Economic Research.
- Kim, J. H. (2014). Identifying the distribution of treatment effects under support restrictions. arXiv preprint arXiv:1410.5885.
- Kreinovich, V. and Ferson, S. (2006). Computing best-possible bounds for the distribution of a sum of several variables is np-hard. *International Journal of Approximate Reasoning*, 41(3):331–342.
- Lu, J., Ding, P., and Dasgupta, T. (2018). Treatment effects on ordinal outcomes: Causal estimands and sharp bounds. *Journal of Educational and Behavioral Statistics*, 43(5):540–567.
- Makarov, G. (1982). Estimates for the distribution function of a sum of two random variables when the marginal distributions are fixed. *Theory of Probability & its Applications*, 26(4):803–806.
- Mullahy, J. (2018). Individual results may vary: Inequality-probability bounds for some healthoutcome treatment effects. *Journal of Health Economics*, 61:151–162.
- Nelsen, R. B. (2006). An introduction to copulas. Springer.
- Rubin, D. B. (1978). Bayesian inference for causal effects: The role of randomization. *The Annals of statistics*, pages 34–58.
- Schmidt, T. (2007). Coping with copulas. Copulas-From theory to application in finance, 3:34.
- Sklar, M. (1959). Fonctions de repartition an dimensions et leurs marges. Publ. inst. statist. univ. Paris, 8:229–231.

Williamson, R. C. and Downs, T. (1990). Probabilistic arithmetic. i. numerical methods for calculating convolutions and dependency bounds. *International journal of approximate reasoning*, 4(2):89–158.